

## SHORT COMMUNICATION

# Genetic Structure of Cactus Pear (*Opuntia Ficus Indica*) in Moroccan Collection

Aissam El Finti, Mohamed Belayadi, Rachida El Boullani, Fouad Msanda, Mohammed Amine Serghini, and Abdelhamid El Mousadik\*

Laboratory of Biotechnologies and Valorization of Natural Resources, University of Ibn Zohr, Faculty of Sciences B.P. 8106, Agadir, Morocco

Received: September 12, 2012 / Accepted: December 18, 2012

## Abstract

Recently, a large plantation has been established in Morocco, to reduce water and wind erosion, rangeland degradation, sand movement and to enhance the restoration of the vegetation cover. However, this plant material has unknown genetic characterization. In addition, several local classifications and morphological descriptions were used. The objective of this study was to analyze the genetic diversity using RAPD markers in a collection of 13 provenances of Moroccan *Opuntia ficus indica* (L.). Based on 13 random primers, the result showed that the level of diversity ( $h$ ) and polymorphism varied according to the provenance. A high genetic differentiation was found between the provenances ( $G_{st} = 0.29$ ), thus some loci were characteristic of certain provenances. These results can be used to characterize genetic resources of Morocco cactus pear, and to initiate a program of genetic improvement and selection.

**Keywords:** *Opuntia ficus indica*, RAPD, Genetic diversity, Genetic resources.

## Introduction

The genus *Opuntia* belongs to the Cactaceae family (subfamily *Opuntioideae*) and includes several species that originated in North and South America, some of which were relatively recently distributed throughout the world. The number of species included in the genus is unknown (Chávez-Moreno et al., 2009). The cactus was introduced to the Mediterranean during the 16<sup>th</sup> century (Barbera, 1995). In Morocco, *Opuntia ficus* represented particularly in arid and semi-arid areas, from Oriental (Oujda, Figuig) to South (Sidi Ifni) through the arid plateaus of Haouz. Cactus pear constitutes, in these areas a resource to control water and wind erosion in eroded soils. The taxonomy of cacti has been traditionally based on comparative observation of morphological and biogeographic data (Metzing and Kiesling, 2008). A detailed description of the germplasm is needed because it may have valuable genes for use in breeding programs.

Due to the multiple uses and the ability of cacti to thrive in arid and semiarid environments, it has become increasingly important to describe and characterize these valuable resources. The latter is a challenging goal since up to now; knowledge regarding the amount of genetic variation and genetic relationship by means of molecular tools is missing in Moroccan *Opuntia ficus indica*. In fact, though this crop is widely cultivated in the country, the majority of research works were especially oriented towards the characterization of the nutritional value of the cladodes as an important fodder crop in arid areas, independently

\* Corresponding author: elmousadik@yahoo.fr

from their genetic potential, (Boujghagh and Chajia, 2001). At present, molecular markers have been proved to be valuable tools in the characterization and evaluation of genetic diversity within and between species and populations. It has been shown that different markers might reveal different classes of variation (Russell et al., 1997). The advent of the polymerase chain reaction (PCR) favored the development of different molecular techniques (Saiki et al., 1998). These molecular markers had been successfully used in *Opuntia* genus for detecting genetic diversity and relationships (Arnholdt-Schmitt et al., 2001; Labra et al., 2003). Of these techniques, RAPD has several advantages, such as simplicity of use, low cost, and the use of small amount of plant material, etc.

The success of cactus plantations in arid and sub-arid environments insists on leading researches on the management of Moroccan genetic resources, which will permit a good knowledge of the planted cacti. At the present, no study is yet available based on genetic variation and relationship by molecular tools in *Opuntia ficus indica* of Morocco.

The objectives of this study were to assess the usefulness of RAPD to differentiate *Opuntia* ecotypes and to investigate genetic relationships among different ecotypes, and to set up rational decisions concerning the establishment of a national reference collection. Indeed, though this crop is widely cultivated in the country, collection repositories are missing.

## Materials and Methods

The Cladodes of *Opuntia ficus indica*, representing 13 provenances were collected from 13 localities in Morocco (Table 1) and planted in Faculty of Sciences Agadir, this material were used for molecular polymorphism (RAPD) research. For each accession, an external slice of the cladode was taken for analysis. A piece of about 1 g of the chlorenchyma was cut using a scalpel and taking care not to include areole. The protocol of DNA extraction used here is that of (Saghai-Marooof et al., 1984) modified. DNA was quantified by visual comparison with lambda DNA molecular marker on ethidium bromide stained agarose gels.

The amplification is performed according to the protocol

(Arnholdt-Schmitt et al., 2001), 14 oligonucleotide (decamer) were used for the amplification of random DNA (OPA-03, OPA-08, OPA-09, OPA-10, OPA-11 OPA-12 OPA-13 OPA-14 OPA-15 OPA-16 OPA-17 OPA-18 OPA-19 OPA-20). PCR reactions were performed in a 25 ml reaction mixture containing: 2.5 ng of template DNA, 2.5 ml of Go Taq buffer (Promega), 3.5 mL of dNTPs, 2 ml of 25 mM MgCl<sub>2</sub>, 25 pmol of primer and 0.5 U of Go Taq DNA polymerase (Promega). The PCR was performed in a Thermoblock thermocycler (Techne).

PCR amplification was performed using the following profile: 5 min at 95°C; 45 cycles of denaturation (1min at 95°C), annealing (1min at 36°C) and extension (5 min at 72°C); then a final step for 5 min at 72°C.

Products of the PCR were separated by electrophoresis in 2% agarose gels using a volt range of 2 Vcm<sup>-1</sup> for 3h. The gel was finally fixed with ethidium bromide and photographed under UV light. Size of alleles was defined using the molecular weight marker 100 bp DNA leaders.

Different RAPD amplified bands were scored for their presence (1) or absence (0) and the resulting data matrices were computed with Nei coefficient to provide a similarity matrix. Different parameters have been estimated and the ability of primers to distinguish between individuals was calculated by evaluating the resolving power, POPGENE32 software was used to treat the binary matrices obtained to estimate genetic parameters intra-provenances (%P: percentage of loci polymorphic, h: Nei's genetic diversity (Nei, 1973) and inter-provenance (Gst: coefficient of genetic differentiation and genetic distance of (Nei, 1978).

## Results and Discussion

Photometric measurements at 260 and 280 nm of the nucleic acid extracts of the various accessions of Barbary fig indicated a quotient (260/280) between 1.82 and 2.02. Thus, the obtained DNA is of high quality.


Among the 14 primers used to assess polymorphism in the tested ecotypes, 13 have revealed unambiguously scorable bands, these mentioned primers generated multiple banding each produced two to twenty readable tapes and are repro-

**Table 1.** Geographic localization of several provenances of *Opuntia ficus indica* under study.


Provenance	Vernacular names	Latitude W	Longitude N	Altitude (m)
Ait Baha	Achefri	30°10'	9°14'	222
Tafraout	Achefri	29°53'	9°00'	1215
Sidi-Ifni M	Moussa	29°20'	10°08'	66
Sidi-Ifni A	Aissa	29°20'	10°08'	65
Tamri	Achefri	30°40'	9°52'	36
Ben Guerir	Aknari	32°30'	7°53'	434
Chaoun	Zaaboul	35°08'	5°16'	555
Hoceima	Delahia	30°57'	4°17'	1100
Nador	Hndiya	35°15'	3°40'	70
Berkane	Hndiya	34°51'	2°36'	200
Meknès	Hndiya	33°47'	5°29'	700
Mohammedia	Hndiya	33°41'	7°19'	64
Safi	Aknari	32°40'	9°04'	56

**Table 2.** Parameters of genetic diversity evaluated on the RAPD markers in Moroccan cacti.

Primers	Number of bands	Number of polymorphic bands	% of polymorphic bands	Number of RAPD profiles	Resolving Power (Rp)
OPA-03	4	1	25%	1	0.24
OPA-08	4	3	75%	4	1.66
OPA-09	2	1	50%	2	0.04
OPA-10	9	4	44.44%	6	1.24
OPA-11	9	8	88.88%	14	2.98
OPA-12	7	5	71.42%	7	2.20
OPA-13	-	-	-	-	-
OPA-14	20	18	90%	22	7.68
OPA-15	13	2	15.38%	4	1.02
OPA-16	6	1	16.66%	2	0.24
OPA-17	7	2	28.57%	3	0.76
OPA-18	7	0	0%	1	0.00
OPA-19	10	4	40%	5	2.10
OPA-20	7	1	14.28%	2	0.10
Total	105	50	-	73	20.26
Means	8.07	3.84	-	-	1.55



**Figure 1.** Nei's Genetic diversity (h) within provenances.



**Figure 2.** Dendrogramme of 13 provenances of *Opuntia ficus indica* throughout Nei's genetic distance.

ducible (Table 2). OPA-13 primer was abandoned because of the difficulty of reading the gel. A total of 105 bands were obtained which 55 were monomorphic and 50 were polymorphic. The average number of bands per primer was 8.07. All bands reported 73 different electrophoretic profiles. The resolving power of primers varied from 0 for the primer OPA-18 to 7.68 for primer OPA-14 with an average value of 1.55. The primer OPA-14, followed by primer OPA-11 and OPA-12, primers seem to be the most efficient to assess the genetic diversity, since they presented relatively high resolving power rates among the prickly pear in Morocco.

Genetic diversity ( $h$ ) has a nil value for the accessions analyzed of Meknes Mohammedia and Safi and a value ranges from 0.01 to 0.11 for other provenances (Fig. 1). Thus, the ecotypes tested from Meknes, Mohammedia and Safi are monomorphic, while other ecotypes have different levels of polymorphism. The ecotype of Chaoun is the most highly polymorphic ( $h = 0.11$ ).

$G_{st}$  coefficient has a value of 0.29 if we takes into account all the loci studied, and 0.75 if we limit only to polymorphic loci. It seems that in both cases, studied provenances have a significant structuring and gene flows are very limited.  $G_{st}$  furthermore the  $G_{st}$  coefficient has the greatest value for loci OPA-11-01-03 and OPA-04 ( $G_{st} = 1$ ). Thus, these loci characterize provenances of Ait Baha and Taфраout. The genetic distance of (Nei, 1978) in terms of dendrogram (Fig. 2 and 3) shows different levels of similarities and dissimilarities between studied provenances. Therefore, apart from provenances of Safi and Mohammedia which are similar, all others provenances shows different cases, including the isolation of Ait Baha and Taфраout on the one hand, and Berkane and Meknes on the other hand. These results showed a disparity between the varieties named Moussa (late) and Aissa (early) from Sidi Ifni. Also, the geographical distribution of the studied provenances is partially respected throughout

the genetic similarities.

## Conclusion

The results presented in this work are the first contributions to the application of molecular markers to assess the genetic diversity of prickly pear in Morocco. Exploring RAPD showed that provenances studied are genetically quite divergent. The loci analyzed participated differently to the description of this divergence. Thus, some loci were polymorphic and easily characterized some provenances.

These results may contribute to the establishment of a conservation program and management of the genetic diversity of prickly pear in Morocco. They can be developed by analyzing other markers with a larger sample. This work could also be combined to agro morphological traits to start a breeding program and a genetic selection.

## References

- Arnholdt-Schmitt B, LC Girao, RM Llamoca-Zarate, and FAP Campos (2001) Genome characterization of *Opuntia ficus-indica*: A simple and efficient micromethod. *J. PACD* 4: 57-65.
- Barbera G (1995) History, Economic and Agro-Ecological Importance. In: *Agroecology, Cultivation and uses of Cactus Pear*, Barbera G, P Inglese and E Pimienta-Barrios (Eds.). FAO Plant Production, Protection Paper, Rome, Italy, pp: 1-2.
- Chávez-Moreno CK, EA Tecante, and EA Casas (2009) The *Opuntia* (Cactaceae) and *Dactylopius* (Hemiptera:Dactylopiidae) in Mexico: a historical perspective of use, interaction and distribution. *Biodivers Conserv* 18: 3337-3355.
- Labra M, F Grassi, M Bardini, S Imazio, and A Guiggi (2003) Genetic relationships in *Opuntia* mill. Genus (cactaceae) detected by molecular marker. *Plant Sci.*, 165(5): 1129-1136.
- Metzing D and R Kiesling (2008) The study of the cactus evolution: the

- pre-DNA era. *Heseltonia* 14: 6–25.
- Boujghagh M and L Chajja (2001) Le cactus: outil de gestion de la sécheresse dans le Sud Marocain. In “Terre et Vie”, Vol. 52, pp. 1-7.
- Nei M (1973) Analysis of gene diversity in subdivided populations. *PNAS USA* 70(12): 3321-332.
- Nei M (1978) Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* 89(3): 583-590.
- Russell JR, JD Fuller, M Macaulay, BG Hatz, A Jahoor, W Powell, and R Waugh (1997) Direct comparison of levels of genetic variation among barley accessions detected by RFLPs, AFLPs, SSRs, and RAPDs. *TAG Theoretical and Applied Genetics* 95(4): 714-722.
- Saghai-Maroo M.A, KM Soliman, RA Jorgensen, and RW Allard (1984) Ribosomal DNA spacer length polymorphism in barley: Mendelian inheritance, chromosomal location and population dynamics. *PNAS* 81(24): 8014-8018.
- Saiki R, DH Gelfand, S Stoffel, SJ Scharf, and R Higuchi (1998) Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* 239(4839): 487-491.