

Plant Growth Promoting Rhizobacteria (PGPR) Isolated from an Arid Soil in Saudi Arabia Improve Maize Growth

Abdelazize Eljiati^{1*}, Yassine Elmaati^{2,3}, and Hammou Ouchaou²

¹ National Centre for Palms & Dates, Al Dahi, Hittin, Riyadh 13512. Saudi Arabia; ² R&D Department. YALA Laboratory. Yousef Abdul Latif and Sons Agriculture Ltd. (YALA) Company, Qassim, Saudi Arabia; ³ Laboratory of Biotechnology and Valorization of Natural Resources, Faculty of Sciences, Ibn Zohr University, Agadir, Morocco

Received: February 19, 2024 / Accepted: April 29, 2024

Abstract

The rhizosphere represents the main source of bacteria commonly referred to as rhizobacteria. Such beneficial rhizobacteria with plant-beneficial activities are generally defined as plant growth promoting rhizobacteria (PGPR). The aim of this study was to investigate the ability of native rhizobacteria (PGPR) isolated from an arid soil of date palm in Al-Qassim region, Saudi Arabia, to enhance plants growth. Maize (*Zea mays* L.) was used as model crop for this research. Maize seedlings roots were inoculated with *Bacillus* and *Enterobacter* bacteria. The seedlings showed significant increases in stem, leaf, and root growth. The maximal shoot lengths were obtained with strain I2: *Bacillus cereus* (95.41 cm) with an increase of 33.45 % compared to uninoculated control seedlings. The three isolates I2: *Bacillus cereus*, AZS2: *Bacillus subtilis* and commercial strain AZB: *Azospirillum brasilense* caused a highly significant increase in the total number of leaves ranging from 10.9% to 12.7% compared to the uninoculated controls. Seedlings inoculated with AZS2: *Bacillus subtilis* strain exhibited the highest aerial dry biomasses with an improvement of more than 85 % (30.76 g) compared with uninoculated control plants and more than 62 % compared to uninoculated NaCl control plants. The inoculation treatment with I2: *Bacillus cereus* strain induced an improvement of more than 65 % (27.44 g) over uninoculated control and more than 45 % over uninoculated NaCl control. The strain

AZS2: *Bacillus subtilis* produced the highest root dry weights, in comparison to other isolates and induced an improvement of 30.17% (26.06 g) compared to uninoculated control plants and 24.09% compared to uninoculated plants (NaCl control). The most effective rhizobacterial treatment in the dry biomasses of whole seedling (aerial dry biomass and root dry biomass) is AZS2: *Bacillus subtilis* strain which induced an improvement of 55% (56.83 g) compared to uninoculated plants (control) and 42% compared to uninoculated plants (NaCl control). The most important production of kernels was recorded with AZS2: *Bacillus subtilis* strain. Therefore, these findings suggested that the use of PGPR strains as inoculant biofertilizers might be beneficial for crop production cultivation especially in arid and semi-arid regions.

Keywords: Plant growth promoting rhizobacteria (PGPR), *Bacillus subtilis*, *Bacillus cereus*, *Enterobacter ludwigii*, inoculation, *Zea mays* L.

* Corresponding author: a.eljiati@gmail.com

Introduction

Bacteria that can aggressively colonize the rhizosphere or plant roots or both and promote growth and yield of plants are referred to as plant growth promoting rhizobacteria (PGPR) (Kloepper et al., 1989). These rhizosphere bacteria enhance crop growth and yield directly, either by promoting nutrition, for example, by phosphate (Hayat et al., 2010; Sharma et al., 2007; Das et al., 2003), and potassium solubilization (Wang et al., 2020; Han et al., 2006) and ammonia production (Mukhtar et al., 2020; Ahmad et al., 2008; Joseph et al., 2007) or by synthesizing metabolites with great agricultural interest such as plant growth regulators and siderophores (Tian et al., 2009; Arkhipova et al., 2005; Barazani et Friedman, 2001). They can also promote growth indirectly, acting as bio-controlling agents for suppression of growth of soil borne phytopathogen microorganisms and as stimulator of other beneficial organisms for the plant (Abbasi et al., 2011; Bhattacharyya and Jha, 2011). PGPR improve soil structure and bioremediate polluted soils by sequestering toxic heavy metal and degrading xenobiotic compounds (Ahmad et al., 2012; Braud et al., 2009). Depending on their beneficial roles in the rhizosphere, PGPR have been classified as biofertilizers, phytostimulators, rhizoremediators and biopesticides (Martínez-Viveros et al., 2010). Consequently, the application of these beneficial microorganisms as bioinoculants appears as an ecological friendly biotechnological tool (Dimpka et al., 2009) to alleviate detrimental effects of intensive farming practices that are using synthetic fertilizers and pesticides without caring about environmental problems and soil health (Elkoca et al., 2010). Numerous laboratory, greenhouse and field studies are available on the screening of PGPR for their multiple plant growth promoting activities (Wang et al., 2020; Gouda et al., 2018; Oteino et al., 2015; Hayat et al., 2010; Joseph et al., 2007) and utilization of PGPR-based products in agricultural crop production systems (Yadav et al., 2017; Cakmakci et al., 2006). These products are mainly applied as seed treatment, soil amendment, or soil drench at the time of sowing or immediately after transplantation, to facilitate better nutrient uptake, greater production of growth hormone and beneficial phytochemicals in crops leading to higher crops yield and quality (Kloepper et al., 2004). PGPR activity has been reported in strains belonging to a several genera, such as *Pseudomonas*, *Azospirillum*, *Azotobacter*, *Klebsiella*, *Enterobacter*, *Rhizobium*, *Bradyrhizobium*, *Alcaligenes*, *Arthobacter*, *Burkholderia*, *Bacillus*, *Serratia* and *Xanthomonas* (Verma et al., 2013; Karnwal 2009; Patten and Glick 1996; Glick, 1995; Kloepper et al., 1989). *Pseudomonas* and *Bacillus* spp. have been the most studied bacteria for their plant growth promotion (PGP) activity and ability to produce beneficial substances (Kejela et al., 2016; Pham et al., 2017). Isolation of native strains adapted to the arid environment may contribute to formulation of inoculants suitable for use in local crops, as they are adapted to the environment and can be, thereby more competent than imported microbial strains. The positive impact of PGPR has been studied in annual crops like wheat (Bashan, 1986), soybeans (Cattelan et al., 1999),

beans (Jarak et al., 2012) and corn (Di Salvo et al., 2018; Ullah et al., 2014) in several ways.

Therefore, this study was designed to select effective strains from a series of native rhizobacteria (PGPR) isolated in an arid area soil of date palm in Al-Qassim region, Saudi Arabia, by maize (*Zea mays* L.) growth promotion assay under greenhouse conditions. These native strains are also compared with a commercial microbial strain, as a positive control. Therefore, the application of the selected effective PGPR strains as microbial inoculants for crops would significantly promote their sustainable production in arid conditions and reduce the use of inorganic fertilizers and pesticides, which often pollute the environment.

Materials and Methods

Bacterial Inoculants

The rhizobacterial strains (I2: *Bacillus cereus*, AZS2: *Bacillus subtilis*, AZA2: *Enterobacter ludwigii* and PSA1: *Enterobacter ludwigii*) used in this study were previously isolated from an arid area soil in Qassim province, Saudi Arabia (Elmaati et al., 2020). These bacterial strains were characterized and selected based on their plant growth promoting traits, comprising very good phosphate and potassium solubilization and ammonium production. A commercial strain (AZB; *Azospirillum brasilense*) was used as a positive control to compare it with these native strains.

Inoculation of Maize Plants by PGPR Strains

To prepare the inoculum for each rhizobacterial strain, pure cultures were grown in nutrient agar (R2A Agar medium). After 48 hours of incubation at 25 °C, a single colony from each strain was transferred into a 200 ml sterilized Erlenmeyer flask, containing sterilized 102 medium broth (= LMG 1089 medium) with the following composition (g L⁻¹): Sucrose, 20.0; Casein hydrolyzate, 16.0; Yeast extract, 8.0; KH₂PO₄, 4.0; MgSO₄ × 7 H₂O, 0.30, and grown aerobically for 4-5 days on a rotating shaker (150 rpm) at 32 °C, to obtain a final concentration of 10⁹ CFU ml⁻¹. After incubation, bacterial growth is estimated by measuring the absorbance of the culture at 600 nm. To wash the bacteria, bacterial cells were centrifuged at 3000 rpm for 10 min in 15 ml tubes. The supernatant was then discarded and the pellet was washed once with 5 ml of sterile NaCl solution (Physiologic Sterile Water (0.85%)) and finally resuspended in 200 ml of the same solution.

Maize (*Zea mays* L.) was used as the test plant for the inoculation in this experiment. Seeds of a homogeneous variety were surface sterilized to eliminate all kinds of contamination according to the method of Götz et al. (2006): The seeds were immersed for 1 min in ethanol (70%) with gentle agitation. They are then put back into 12% diluted sodium hypochlorite solution containing three drops of wetting agent (Tween 20) for 15 minutes. To get rid of the chlorine, the seeds were rinsed several times with sterile distilled water.

Sterilized seeds were sown in alveolar plates containing

autoclaved (1:1 v/v) mixture of peat and vermiculite respectively. They were then placed in a greenhouse at a natural photoperiod (at a temperature of 28 to 40 °C) and are regularly irrigated.

Twenty days after germination, the roots of maize seedlings were dipped into the inocula for 2 h at 25 °C. Control seedlings were divided into two groups; the first is dipped into sterile NaCl solution as the bacteria washing procedure containing this solution, while the second is dipped into sterile distilled water. Each treated seedling was planted in a disinfected and labeled plastic pot (8 L) containing autoclaved (1/1: v/v) mixture of soil and vermiculite respectively. Pots were then placed in a greenhouse at a natural photoperiod (at a temperature of 28 to 40 °C). The experiment was carried out in a completely randomized block design with 5 replications for each treatment. Seedlings were irrigated with 300 ml of well water every two days to maintain at field capacity and received no fertilizers.

Estimation of Some Agro Morphological Parameters of Plants

To evaluate the response to rhizobacterial inoculation, during the experiment, a daily monitoring of the evolution of the growth of the maize plants was done after the application of the inoculum for a period of 90 days. The height from the collar (size of the aerial part) and the number of leaves are the main growth parameters used in this study. These two growth parameters were measured at the start and every ten days during the 3 months of treatment. Plant height was determined by measuring from the plant's base to the top of the newest fully developed leaf.

At the end of the experiment, the shoots and roots of each plant were put in paper bags and dried in an oven at 65°C for 72 hours (Sfairi, 2013) to report the total dry biomass (shoot and root dry biomass).

Statistical Data Analysis

The mean value of each treatment, as well as the corresponding standard deviation, were calculated using the data of all the replicates carried out. The data obtained was analyzed statistically using R 3.2.0 and multivariate analyses were performed using R language (Dray & Dufour, 2007; R Development Core Team, 2011).

Results

Effect of Rhizobacterial Inoculation on Shoot Length

The rhizobacterial isolates (AZB, AZS2, I2, PSA1, and AZA2) significantly affected the shoot length of maize seedlings. Results reveal that PGPR promoted an increase in shoot length over un-inoculated (control) (Figure 1 - A and B). The maximal lengths of the maize seedlings were obtained with strain (I2) (95.41 cm) with an increase of 33.45 % compared to un-inoculated (controls). Treatment of maize seedlings with strain (I2) showed a significant increase in shoot length improvement

rates compared to other strains including the commercial strain (AZB). There was no significant difference in shoot length between the two controls.

Effect of Rhizobacterial Inoculation on Leaf Numbers

From 1 to 38 days after emergence (Figure 2A), the number of leaves of maize plants increased linearly in all treatments. These results were identical for all the rhizobacterial isolates and the controls. However, we noted the beginning of the stability of the number of leaves between the 40th and the last days of the cycle in some plants.

The effects of different rhizobacterial isolates were significant on the total number of leaves of maize, compared with control. Application of AZS2 isolate to maize seedlings recorded non-significantly higher number of leaves, compared with AZB and I2, and all these three isolates were comparatively more effective than rest of the isolates and the two un-inoculated control. The three effective isolates caused a highly significant increase in the total number of leaves ranging from 10.9% to 12.7% compared to the un-inoculated control. No significant difference was observed between the two isolates (PSA1 and AZA2) and the control with NaCl (Figure 2B).




Figure 1. (A) Effect of rhizobacterial inoculation on shoot length (B) Height growth of maize seedlings after 50 days of cultivation under the effect of applied rhizobacterial strains.

Effect of Rhizobacterial Inoculation on Leaf Desiccation

During the period of evolution of the different foliar levels of the maize seedlings, we noticed that the leaves of the base wither and turn brown, by a yellowing and drying which begins with the end of the leaves and extends thereafter, until they dry out completely. The number of dried older leaves was recorded for each plant. At the end of the experiment, the results of the analysis of variance relating to this parameter show that there is no statistically significant difference between the rhizobacteria and the controls without bacteria (Figure 3).

Effect of Rhizobacterial Inoculation on Aerial Dry Biomass

The analysis of variance relative to the aerial dry biomasses of maize seedling shows that there is a highly significant difference between the different strains and the un-inoculated control and un-inoculated control NaCl control (with a total aerial dry biomass which marked a rate of 16.59 g and 18.88 g respectively). Seedlings inoculated with AZS2 strain exhibited the highest aerial dry biomasses with an improvement of more than 85 % (30.76 g) compared with un-inoculated control plants and more than 62 % compared to un-inoculated NaCl control plants. The inoculation treatment with I2 strain




Figure 2. (A) Evolutionary trend of the adjusted mean of the number of maize leaves (B) Effect of rhizobacterial inoculation on leaf numbers.




Figure 3. Effect of rhizobacterial inoculation on leaf desiccation.




Figure 4. Effect of rhizobacterial inoculation on aerial dry biomass

induced an improvement of more than 65 % (27.44g) over un-inoculated control and more than 45 % over un-inoculated NaCl control. On the other hand, the commercial strain (AZB) showed an improvement of 64.98% (27.37 g) compared to untreated control and 44.97% compared to untreated NaCl control. The others two strains (PSA1 and AZA2) are characterized by their lowest significant effect on aerial dry biomass per comparison to the both control plants (control and NaCl control) (up to 38 % increase) (Figure 4).

Effect of Rhizobacterial Inoculation on Root Dry Biomass

The effect of the five rhizobacterial strains on the root dry biomasses of maize plants is illustrated in (Figure 5A). All the tested strains were significantly improved the root dry biomasses in comparison with controls (control and NaCl control), which scored a rate of 20.02 g and 21.00 g respectively. The strain AZS2 produced the highest root dry weights, in comparison to other isolates and induced an improvement of 30.17% (26.06 g) compared to un-inoculated plants (control) and 24.09% compared to un-inoculated plants (NaCl control). While, in comparison with controls, the inoculation with strains I2, PSA1 and AZB showed a significant difference with improvement rates ranging from 16.2% to 19.35%.

It is observed from the results (Figure 5B) that the rhizobacterial strains caused greater increase in root system of maize plants as compared with controls (control and NaCl control). Indeed, the highest root lengths was recorded with the inoculation of AZS2 and I2 strains), in comparison to other strains.

Effect of Rhizobacterial Inoculation on Whole Seedling Dry Biomass

A very highly significant improvement rate in the dry biomasses of whole seedling (aerial dry biomass and root dry biomass) inoculated with the rhizobacterial strains is recorded compared to the un-inoculated plants (control and NaCl control), which scored a level of 36.62 g and 39.94 g respectively. The most effective rhizobacterial treatment is AZS2 strain which induced an improvement of 55% (56.83 g) compared to un-inoculated plants (control) and 42% compared to un-inoculated plants (NaCl control). Analysis of the variance applied to dry biomasses of whole seedling indicated that there was no significant difference between other strains (I2, AZA2, PSA1 and AZB), but they induced an increase for this parameter compared to the un-inoculated control (Figure 6).




Figure 5. (A) Effect of rhizobacterial inoculation on root dry biomass (B) Effect of rhizobacterial inoculation on root length of maize seedlings

Effect of Rhizobacterial Inoculation on Kernel Numbers

All the inoculated treatments proved statistically superior over un-inoculated control in improving number of kernels. No significant difference was observed between the four bacterial strains AZB, AZS2, I2 and PSA1. However, the most important production of kernels was recorded with AZS2 strain. While total kernels number was significantly increased with these four strains as compared to AZA2 strain and compared to un-inoculated control (Figure 7).

Discussion

Plant Growth Promoting Rhizobacteria (PGPR) are free-living microbes that live on or around the roots (Kloepper et al., 1989) and promote plant growth and yield (Wu et al., 2005). Native rhizobacteria (PGPR) isolated in arid area soil of date palm in Al-Qassim region, Saudi Arabia, were used in this study to constitute the rhizobacterial inoculum, which was then used to inoculate maize (*Zea mays* L.) seedlings. These strains exhibited significant plant growth promoting attributes in vitro tests and selected from previous screening experiments (Elmaati 2020). In addition to its nutritional and




Figure 6. Effect of rhizobacterial inoculation on whole seedling dry biomass.




Figure 7. Effect of rhizobacterial inoculation on kernel numbers.

economic importance, maize has been a keystone model organism for basic and applied research in plant biology (Strable and Scanlon, 2009).

Overall, the results of the study of the growth of maize seedlings treated with the various rhizobacterial strains enabled us to conclude that the efficacy of different strains for growth-promoting of maize was variable. Indeed, the investigated PGPR strains in this work showed positive PGP traits. These potentialities seem playing an effective role for the plant in helping it to better absorb nutrients. PGPR have long been known to promote growth when added to seeds, roots or tubers in a wide range of plant species (Kloepper et al., 1980), increasing both growth and yield (Wu et al., 2005) by improving the concentration of nutrients in the host plant (Canbolat et al., 2006).

Several studies have reported that inoculation of maize plants with PGPR strains caused significant increase in plant height, plant dry weight, stem diameter, root length and weight, yield, number of leaves and leaf area, and plant nutrient uptake of N, P, K, Fe, Zn, Mn and Cu (Yazdani et al., 2009; Jarak et al., 2012; Gholami et al., 2012; Calvo et al., 2017). According to our results maize seedlings inoculated by dipping the roots in bacterial suspensions showed a statistically significant improvement in the growth parameters studied as compared to treatments without inoculation. Interestingly, the plant height, number of leaves, shoot dry weight, root dry weight and kernels number were significantly higher with plants treated with AZS2 strain (*Bacillus subtilis*) followed by I2 strain (*Bacillus cereus*) in comparison with un-inoculated control and with other strains including the commercial strain (AZB, *Azospirillum brasilense*). This is consistent with previous studies, which demonstrated that plant growth-promoting activities of *Bacillus* spp. are well characterized as evidenced by increased growth of roots, shoots, and leaves as well as enhanced yields. In this context, increased plant height and shoot biomass of *Arabidopsis*, corn, and tomato under greenhouse conditions have been reported by inoculating with four isolated *Bacillus* strains from rainforest soils (Huang et al., 2015). Results obtained by Hassan (2017) report that *B. cereus* Tp.1B and *B. subtilis* Tp.6B strains significantly increased root length and root weight in maize compared to controls. Co-inoculation of *Bacillus* spp with other PGPR strains reduces phosphorus demand by 50% without affecting maize yield (Yazdani et al., 2009). Moreover, Ferreira et al., 2018 reported that *Bacillus subtilis* promotes positive influence on plant growth of maize plants under normal conditions (without salinity). *Bacillus subtilis* strain was the most effective in promoting nitrogen accumulation and, therefore, increased chlorophyll content in maize (Aquino et al., 2019; Almaghrabi et al., 2014). When tomato seeds were treated with *Bacillus subtilis* (EPC016), a significant increase in seedling growth was observed relative to un-inoculated plants (Ramayabharathi et al., 2013). In another study, Tilak and Reddy (2006) observed a significant increase in grain yield rate of 43.8% in maize plants inoculated with *Bacillus cereus*. This last strain was found to exhibit the highest nitrogenase activity among 42 different strains of *Bacillus* spp studied by Am-

brocini et al., (2016). In addition, *B. cereus* and *B. megaterium* have been reported as organic phosphorus mineralizing bacteria (Guang Can et al., 2008). The works of Habib et al., (2015, 2016) on rhizobacteria isolated from saline soil and selected for their PGP activities revealed that they showed significant salt tolerance properties. These rhizobacteria were identified as *Enterobacter* sp. and *Bacillus cereus*.

Our research indicates that significantly lower values of different growth parameters were recorded in maize plants inoculated with *Enterobacter ludwigii* in comparison to *Bacillus subtilis* and *Bacillus cereus*. On the other hand, and compared to the un-inoculated control plants, *Enterobacter ludwigii* had significant positive effects on maize plant growth parameters. Zaballa et al., (2020) found that barley plants inoculated with the *Enterobacter ludwigii* strain showed improvement in growth and phosphate uptake compared to the un-inoculated control. Tahir et al., (2013) reported that inoculation of wheat plants with phosphate-solubilizing and phytohormone-producing bacterial strains such as *Azospirillum*, *Bacillus* and *Enterobacter* improved growth and yield. Moreover, several studies have demonstrated the effectiveness of inoculating wheat grains with different rhizobacteria on plant growth (Abbasi et al., 2011; Rana et al., 2011; Banerjee et al., 2010). Numerous studies have highlighted the increase in dry matter weight of aerial parts in wheat (Bashan, 1986) and maize (García de Salamone and Döbereiner, 1996; Ullah et al., 2014; Di Salvo et al., 2018). The positive effects of PGPR on the yield and growth of crops such as wheat (Ozturk et al., 2003; Salanture et al., 2006) maize (Egamberdiyeva, 2007; Ullah S and B Asghari, 2015; Pereira et al., 2020) soybean (Cattelan et al., 1999) and sugar beet (Cakmakc et al., 2006) have been explained by the ability of these PGPR to fix N₂, solubilize phosphate and produce phytohormones. Thus, these rhizobacteria can be considered as an excellent tool for increasing the availability of phosphorus in plants by mineralization of soil organic phosphorus and by solubilization of phosphate precipitates (Kucey et al., 1989; Pradhan and Sukla, 2006), production of AIA (Chaiharn and Lumyong, 2011; Swain et al., 2007), HCN (Bakker and Schippers, 1987), ammonia (NH₃) (Yadav et al., 2010) and siderophores (Boopathi and Rao, 1999).

In general, our study clearly showed that the inoculation of maize plants with the rhizobacterial strains significantly promoted maize plants growth. These results suggest that these PGPR strains can be applied as biofertilizers for improving plants production. Furthermore, their use can be an ecological alternative to reduce the dependence on chemical fertilizers.

Conclusion

The rhizobacterial strains investigated in our study showed their plant growth ability. These native strains, which belong to the genera *Bacillus* and *Enterobacter*, significantly enhanced the growth of maize plants when compared with the un-inoculated control plants.

The maximal lengths of the maize seedlings were obtained

with strain I2: *Bacillus cereus* (95.41 cm) with an increase of 33.45 % compared to un-inoculated (controls). The three isolates I2: *Bacillus cereus*, AZS2: *Bacillus subtilis* and commercial strain AZB: *Azospirillum brasilense* caused a highly significant increase in the total number of leaves ranging from 10.9% to 12.7% compared to the un-inoculated control. Seedlings inoculated with AZS2: *Bacillus subtilis* strain exhibited the highest aerial dry biomasses with an improvement of more than 85 % (30.76 g) compared with un-inoculated control plants and more than 62 % compared to un-inoculated NaCl control plants. The inoculation treatment with I2: *Bacillus cereus* strain induced an improvement of more than 65 % (27.44g) over un-inoculated control and more than 45 % over un-inoculated NaCl control. The strain AZS2: *Bacillus subtilis* produced the highest root dry weights, in comparison to other isolates and induced an improvement of 30.17% (26.06 g) compared to un-inoculated plants (control) and 24.09% compared to un-inoculated plants (NaCl control). The most effective rhizobacterial treatment in the dry biomasses of whole seedling (aerial dry biomass and root dry biomass) is AZS2: *Bacillus subtilis* strain which induced an improvement of 55% (56.83 g) compared to un-inoculated plants (control) and 42% compared to un-inoculated plants (NaCl control). The most important production of kernels was recorded with AZS2: *Bacillus subtilis* strain.

Consequently, this finding suggests that these PGPR strains could be useful for the development of inoculants biofertilizers to improve the quality and the health of the soil and the plant species by increasing the nutrient availability for the soil and plants especially in arid and semi-arid regions. Furthermore, using biofertilizers that contain these rhizobacterial strains will led to a decrease in the use of chemical fertilizers and will provide high quality products free of harmful agrochemicals for human and environment.

Acknowledgements

The authors would like to deeply thank Mr Youssef Jameel the Chairman of Yousef Bin Abdul Latif and Sons Agriculture Co. Ltd for his financial support and for allowing us to use the farm and laboratory facilities of his Company during this research.

References

- Abbasi MK, S Sharif, M Kazmi, T Sultan and M Aslam (2011) Isolation of plant growth promoting rhizobacteria from wheat rhizosphere and their effect on improving growth, yield and nutrient uptake of plants. *Plant Biosystems* 145: 159-168. DOI: <https://doi.org/10.1080/11263504.2010.542318>.
- Ahemad M (2012) Implications of bacterial resistance against heavy metals in bioremediation: a review. *Journal of Institute of Integrative Omics and Applied Biotechnology (IIOAB)* 3: 39-46.
- Ahmad F, I Ahmad and MS Khan (2008) Screening of free-living rhizospheric bacteria for their multiple plant growth promoting activities. *Microbiological Research* 163: 173-181. DOI: [10.1016/j.micres.2006.04.001](https://doi.org/10.1016/j.micres.2006.04.001).
- Almaghrabi AO, TS Abdelmoneim, H Albishri and T Moussa (2014) Enhancement of maize growth using some plant growth promoting Rhizobacteria (PGPR) under laboratory conditions. *Life Science Journal* 11(11): 764-772.
- Ambrosini A, T Stefanski, BB Lisboa, A Beneduzi, LK Vargas and LMP Passaglia (2016) Diazotrophic bacilli isolated from the sunflower rhizosphere and the potential of *Bacillus mycoides* B38V as bio-fertiliser. *Ann Appl Biol* 168: 93-110.
- Aquino JPA, JFB Macedo, JEL Antunes, MVB Figueiredo, FA Neto and ASF Araujo (2019) Plant growth-promoting endophytic bacteria on maize and sorghum. *Pesquisa Agropecuária Tropical* 49. DOI: [10.1590/1983-40632019v4956241](https://doi.org/10.1590/1983-40632019v4956241).
- Arkhipova TN, SU Veselov, AI Melentiev, EV Martynenkom and GR Kudoyarova (2005) Ability of bacterium *Bacillus subtilis* to produce cytokinins and to influence the growth and endogenous hormone content of lettuce plants. *Plant Soil* 272: 201-209.
- Bakker AW and B Schippers (1987) Microbial cyanide production in the rhizosphere to potato yield reduction and *Pseudomonas* spp. mediated plant growth stimulation. *Soil Biology and Biochemistry* 19: 451-457. DOI: [10.1016/0038-0717\(87\)90037-X](https://doi.org/10.1016/0038-0717(87)90037-X).
- Banerjee S, R Palit, C Sengupta and D Standing (2010) Stress induced phosphate solubilization by *Arthrobacter* sp. and *Bacillus* sp. isolated from tomato rhizosphere. *Australian Journal of crop science* 4: 378-389.
- Barazani O and J Friedman (2001) Allelopathic bacteria and their impact on higher plants. *Critical Reviews in Microbiology* 27(1): 41-55. DOI: [10.1080/20014091096693](https://doi.org/10.1080/20014091096693).
- Bashan Y (1986) Alginate beads as synthetic inoculant carriers for the slow release of bacteria that affect plant growth. *Applied and Environmental Microbiology* 51(5):1089-1098. DOI: [10.1128/aem.51.5.1089-1098.1986](https://doi.org/10.1128/aem.51.5.1089-1098.1986).
- Bhattacharyya PN and DK Jha (2012) Plant growth-promoting rhizobacteria (PGPR), emergence in agriculture. *World J Microbiol Biotechnol* 28: 1327-1350. DOI: [10.1007/s11274-011-0979-9](https://doi.org/10.1007/s11274-011-0979-9).
- Boopathi E and KS Rao (1999) A siderophore from *Pseudomonas putida* type A1, structural and biological characterization. *Biochimica et Biophysica Acta (BBA)- Protein structure and molecular enzymology* 1435: 30-40. DOI: [10.1016/S0167-4838\(99\)00204-6](https://doi.org/10.1016/S0167-4838(99)00204-6).
- Braud, A., Jézéquel, K., Bazot, S., Lebeau, T. (2009). Enhanced phyto-extraction of an agricultural Cr-, Hg-and Pb-contaminated soil by bioaugmentation with siderophore producing bacteria. *Chemosphere* 74, 280–286. DOI: [10.1016/j.chemosphere.2008.09.013](https://doi.org/10.1016/j.chemosphere.2008.09.013).
- Cakmakci R, F Donmez, A Aydin and F Sahin (2006) Growth promotion of plants by plant growth-promoting rhizobacteria under greenhouse and two different field soil conditions. *Soil Biol. Biochem* 38, 1482-1487. DOI: [10.1016/j.soilbio.2005.09.019](https://doi.org/10.1016/j.soilbio.2005.09.019).
- Calvo P, DB Watts, JW Kloepper and HA Torbert (2017) Effects of Microbial-Based Inoculants on Nutrient Concentrations and Early Root Morphology of Corn (*Zea mays*). *Journal of Plant Nutrition and Soil Science* 180: 56-70. DOI: [10.1002/jpln.201500616](https://doi.org/10.1002/jpln.201500616).
- Canbolat MY, S Bilen, R Çakmakç, F Şahin and A Aydın (2006) Effect of plant growth-promoting bacteria and soil compaction on barley seedling growth, nutrient uptake, soil properties and rhizosphere microflora. *Biol. Fertil. Soils* 42: 350-357. DOI: [10.1007/s00374-005-0034-9](https://doi.org/10.1007/s00374-005-0034-9).
- Cattelan AJ, PG Hartel and JJ Fuhrmann (1999) Screening for plant growth-promoting rhizobacteria to promote early soybean growth. *Soil Sci. Soc. Am. J* 63:1670-1680. DOI: [10.2136/sssaj1999.6361670x](https://doi.org/10.2136/sssaj1999.6361670x).
- Chaiarn M and S Lumyong (2011) Screening and optimization of indole-3-acetic acid production and phosphate solubilization from rhizobacteria aimed at improving plant growth. *Current microbiology* 62: 173-181. DOI: [10.1007/s00284-010-9674-6](https://doi.org/10.1007/s00284-010-9674-6).

- Das AJ, M Kumar and R Kumar (2013) Plant growth promoting PGPR: an alternative of chemical fertilizer for sustainable environment friendly agriculture. *Res J Agric For Sci* 1:21–23.
- Dimkpa C, T Weinand, F Asch (2009) Plant-rhizobacteria interactions alleviate abiotic stress conditions *Plant Cell Environ* 32: 1682–1694. DOI: 10.1111/j.1365-3040.2009.02028.x.
- Di Salvo LP, GC Cellucci, ME Carlino and IE García de Salamone (2018) Plant growth-promoting rhizobacteria inoculation and nitrogen fertilization increase maize (*Zea mays* L.) grain yield and modified rhizosphere microbial communities. *Applied Soil Ecology* 126: 113–120. DOI: 10.1016/j.apsoil.2018.02.010.
- Dray S and AB Dufour (2007) The ade4 package: implementing the duality diagram for ecologists. *J. Stat. Software* 22: 1–20. DOI: 10.18637/jss.v022.i04.
- Egamberdiyeva D (2007) The effect of plant growth promoting bacteria on growth and nutrient uptake of maize in two different soils. *Applied Soil Ecol* 36:184–189. DOI: 10.1016/j.apsoil.2007.02.005.
- Elkoca E, M Turan and M Figen Donmez (2010) effects of single, dual and triple inoculations with *Bacillus subtilis*, *Bacillus megaterium* and *Rhizobium leguminosarum* bv. *phaseoli* on nodulation, nutrient uptake, yield and yield parameters of common bean (*Phaseolus vulgaris* L.), *Journal of Plant Nutrition*, 33 (14): 2104–2119. DOI: 10.1080/01904167.2010.519084.
- Elmaati Y, F Msanda, A Eljati, H Ouchaou, H Boubaker and M Ait Hamza (2020) Characterization of Plant Growth Promoting Rhizobacteria Isolated from an Arid Area Soil of Date Palm in Saudi Arabia. *Journal of Applied Sciences* 20: 196–207. DOI: 10.3923/jas.2020.196.207.
- Ferreira NC, R de CL Mazzuchelli, AC Pacheco, FF de Araujo, JEL Antunes and ASF de Araujo, (2018). *Bacillus subtilis* improves maize tolerance to salinity. *Ciência Rural* 48 (8). DOI: 10.1590/0103-8478cr20170910.
- García de Salamone IE and J Döbereiner (1996) Maize genotype effects on the response to *Azospirillum* inoculation. *Biol. Fertil. Soils* 21 (3): 193–196.
- Gholami A, A Biyari, M Gholipour and HA Rahmani (2012) Growth Promotion of Maize (*Zea mays* L.) by Plant-Growth-Promoting Rhizobacteria under Field Conditions. *Communications in Soil Science and Plant Analysis* 43: 1263–1272. DOI: 10.1080/00103624.2012.666302.
- Glick BR (1995) The enhancement of plant growth by free living bacteria. *Can. J. Microbiol* 41: 109–114. DOI: 10.1139/m95-015.
- Götz M, H Nirenberg, S Krause, H Wolters, S Draeger, A Buchner, Lottmann, G Berg and K Smalla (2006) Fungal endophytes in potato roots studied by traditional isolation and cultivation-independent DNA-based methods. *FEMS. Microbiol. Ecol* 58: 404–413. DOI: 10.1111/j.1574-6941.2006.00169.x.
- Gouda S, RG Kerry, G Das, S Paramithiotis, HS Shin and JK Patra (2018) Revitalization of plant growth promoting rhizobacteria for sustainable development in agriculture. *Microbiol. Res* 206: 131–140. DOI: 10.1016/j.micres.2017.08.016.
- Guang Can TAO, T Shu Jun, CAI Miao Ying and XIE Guang Hui (2008) Phosphate solubilizing and mineralizing abilities of bacteria isolated from soils. *Pedosphere* 18: 515–523. DOI: 10.1016/S1002-0160(08)60042-9.
- Habib SH, H Kausar and HM Saud (2016) Plant growth-promoting Rhizobacteria enhance salinity stress tolerance in Okra through ROS-scavenging enzymes. *Bio Med research international* 2016: 10. DOI: 10.1155/2016/6284547.
- Habib SH, H Kausar, HM Saud, MR Ismail and R Othman (2015) Molecular characterization of stress tolerant plant growth promoting rhizobacteria (PGPR) for growth enhancement of rice. *International Journal of Agriculture and Biology* 18 (01): 184–191. DOI: 10.17957/ijab/15.0094.
- Han, HS, Supanjani and KD Lee (2006) Effect of co-inoculation with phosphate and potassium solubilizing bacteria on mineral uptake and growth of pepper and cucumber. *Plant Soil and Environment* 52: 130–136.
- Hassan SED (2017) Plant growth-promoting activities for bacterial and fungal endophytes isolated from medicinal plant of *Teucrium polium* L. *Journal of Advanced Research* 8 (6): 687–695. DOI: 10.1016/j.jare.2017.09.001.
- Hayat R, S Ali, U Amara, R Khalid and I Ahmed (2010) Soil beneficial bacteria and their role in plant growth promotion: A review. *Anal. Microbiol* 60: 579–598. <https://doi.org/10.1007/s13213-010-0117-1>.
- Huang X, D Zhou, J Guo, DK Manter, KF Reardon and JM Vivanco (2015) *Bacillus* spp. from Rainforest Soil Promote Plant Growth under Limited Nitrogen Conditions. *Journal of Applied Microbiology* 118 (3): 672–684. DOI: 10.1111/jam.12720.
- Jarak M, N Mrkovački, D Bjelić, D Jošić, T Hajnal-Jafari and D Stamenov (2012) Effects of Plant Growth Promoting Rhizobacteria on Maize in Greenhouse and Field Trial. *African Journal of Microbiology Research* 6: 5683–5690. DOI: 10.5897/AJMR12.759.
- Joseph B, RR Patra and R Lawrence (2007) Characterization of plant growth promoting rhizobacteria associated with chickpea (*Cicer arietinum* L.). *Int. J. Plant Prod* 2: 141–152. DOI: 10.22069/ijpp.2012.532.
- Kang, S. M., Radhakrishnan, R., Lee, K. E., You, Y. H., Ko, J. H., Kim, J. H., et al. (2015). Mechanism of plant growth promotion elicited by *Bacillus* sp. LKE15 in oriental melon. *Acta Agric. Scand. Sect. B Soil Plant Sci.* 65, 637–647. DOI: 10.1080/09064710.2015.1040830.
- Karnwal A (2009) Production of indole acetic acid by fluorescent *Pseudomonas* in the presence of l-tryptophan and rice root exudates. *Journal of Plant Pathology, Società Italiana di Patologia Vegetale (SIPaV)*. 91 (1): 61–63. DOI: 10.4454/jpp.v91i1.624.
- Kejela T, VR Thakkar and P Thakor (2016) *Bacillus* species (BT42) isolated from *Coffea arabica* L. rhizosphere antagonizes *Colletotrichum gloeosporioides* and *Fusarium oxysporum* and also exhibits multiple plant growth promoting activity. *BMC Microbiol* 16: 277. DOI: 10.1186/s12866-016-0897-y.
- Kloepper JW, CM Ryn and S Zhang (2004) Induced systemic resistance and promotion of plant growth by *Bacillus* sp. *Phytopathol* 94:1259–1266. DOI: 10.1094/PHYTO.2004.94.11.1259.
- Kloepper JW, J Leong, M Teintze and MN Schroth (1980) Enhanced Plant-Growth by Siderophores Produced by Plant Growth-Promoting Rhizobacteria. *Nature* 286: 885–886. DOI: 10.1038/286885a0.
- Kloepper JW, R Lifshitz and RM Zablutowicz (1989) Free Living Bacterial Inocula for Enhancing Crop Productivity. *Trends in Biotechnology* 7: 39–44. [https://doi.org/10.1016/0167-7799\(89\)90057-7](https://doi.org/10.1016/0167-7799(89)90057-7)
- Kucey RMN, HH Janzen and ME Legget (1989) Microbial mediated increases in plant available phosphorus. *Adv. Agron* 42,199–228. DOI: 10.1016/S0065-2113(08)60525-8.
- Martínez-Viveros O, M Jorquera, DE Crowley, G Gajardo and ML Mora (2010) Mechanisms and practical considerations involved in plant growth promotion by Rhizobacteria. *Journal of Soil Science and Plant Nutrition* 10: 293–319. DOI: 10.4067/S0718-95162010000100006.
- Mukhtar T, S Rehman, D Smith, T Sultan and MF Seleiman et al (2020) Mitigation of heat stress in *Solanum lycopersicum* L. by ACC-deaminase and exopolysaccharide producing *Bacillus cereus*: Effects on biochemical profiling. *Sustainability* 12. <https://doi.org/10.3390/su12062159>.
- Oteino N, RD Lally, S Kiwanuka, A Lloyd, D Ryan, KJ Germaine, and

- DN Dowling (2015) Plant growth promotion induced by phosphate solubilizing endophytic *Pseudomonas* isolates. *Front. Microbiol* 6. DOI: 10.3389/fmicb.2015.00745.
- Ozturk A, O Caglar and F Sahin (2003) Yield response of wheat and barley to inoculation of plant growth promoting rhizobacteria at various levels of nitrogen fertilization. *J. Plant Nutr. Soil Sci* 166: 262-266. DOI:10.1002/JPLN.200390038.
- Patten CL and BR Glick (1996) Bacterial biosynthesis of indole-3-acetic acid. *Can J Microbiol* 42: 207-220. DOI: 10.1139/m96-032.
- Pereira SIA, D Abreu, H Moreira, A Vega, PML Castro (2020) Plant growth-promoting rhizobacteria (PGPR) improve the growth and nutrient use efficiency in maize (*Zea mays* L.) under water deficit conditions. *Heliyon* 6(10). DOI: 10.1016/j.heliyon.2020.e05106.
- Pham TPT, F Cadoret, M Tidjani Alou, S Brah, B Ali Diallo, A Diallo, C Sokhna, J Delerce, PE Fournier, M Million, D Raoult (2017) 'Urmittelita timonensis' gen. nov., sp. nov., 'Blautia marasmi' sp. nov., 'Lachnoclostridium pacaense' sp. nov., 'Bacillus marasmi' sp. nov. and 'Anaerotruncus rubiinfantis' sp. nov., isolated from stool samples of undernourished African children. *New Microbes and New Infections* 17: 84-88. DOI: 10.1016/j.nmni.2017.02.004.
- Pradhan N and LB Sukla (2006) Solubilization of inorganic phosphates by fungi isolated from agriculture soil. *African Journal of Biotechnology* 5: 850-854. <http://www.academicjournals.org/AJB>.
- Radhakrishnan R, Hashem A and Abd_Allah EF (2017) *Bacillus*: A Biological Tool for Crop Improvement through Bio-Molecular Changes in Adverse Environments. *Front. Physiol.* 8:667. DOI: 10.3389/fphys.2017.00667.
- Ramyabharathi S, B Meena and T Raguchander (2013) Induction of defense enzymes and proteins in tomato plants by *Bacillus subtilis* EPCO16 against *Fusarium oxysporum* f. sp. lycopersici. *Madras Agric J.* 100: 126-130.
- Rana A, B Saharan, M Joshi, R Prasanna, K Kumar and L Nain (2011) Identification of multi-trait PGPR isolates and evaluating their potential as inoculants for wheat. *Annals of microbiology* 61: 893-900. DOI: 10.1007/s13213-011-0211-z.
- R Development Core Team, 2011. R: A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing.
- Salanture A, A Ozturk and S Akten (2006) Growth and yield response of spring wheat (*Triticum aestivum* L.) to inoculation with rhizobacteria. *Plant Soil Environ* 52: 111-118. DOI:10.17221/3354-PSE
- Sfairi Y (2013) *Cupressus atlantica* Gaussen, une espèce endémique et menacée du Haut Atlas Occidental Marocain : Etude de sa variabilité morphométrique et chimique et de son comportement écophysiolgique. Université Cadi Ayyad. Faculté des Sciences Semlalia- marrakech. 115 p.
- Sharma K, G Dak, A Agrawal, M Bhatnagar and R Sharma (2007) Effect of phosphate solubilizing bacteria on the germination of *Cicer arietinum* seeds and seedling growth. *Journal of Herbal Medicine and Toxicology* 1: 61-63.
- Strable J and MJ Scanlon (2009) Maize (*Zea mays*): A Model Organism for Basic and Applied Research in Plant Biology. *Cold Spring Harbor Protocols*, pdb.emo132. DOI: 10.1101/pdb.emo132.
- Swain MR, SK Naskar and RC Ray (2007) Indole-3-acetic acid production and effect on sprouting of yam (*Dioscorea rotundata* L.) minisets by *Bacillus subtilis* isolated from culturable cowdung microflora. *Polish Journal of Microbiology* 56:103-117. PMID: 17650680.
- Tahir M, MS Mirza, A Zaheer, MR Dimitrov, H Smidt and S Hameed (2013) Isolation and identification of phosphate solubilizer *Azospirillum*, *Bacillus* and *Enterobacter* strains by 16SrRNA sequence analysis and their effect on growth of wheat (*Triticum aestivum* L.). *Aust. J. Crop Sci* 7: 1284-1292.
- Tian F, Y Ding, H Zhu, L Yao and B Du (2009) Genetic diversity of siderophoreproducing bacteria of tobacco rhizosphere. *Brazilian Journal of Microbiology* 40: 276-284. DOI: 10.1590/S1517-838220090002000013.
- Tilak BR and BS Reddy (2006) *Bacillus cereus* and *B. circulans*-novel inoculants for crops. *Curr Sci* 90: 642-644.
- Ullah S and B Asghari (2015) Isolation of plant-growth-promoting rhizobacteria from rhizospheric soil of halophytes and their impact on maize (*Zea mays* L.) under induced soil salinity. *Canadian Journal of Microbiology*, 61(4): 307-313. DOI: 10.1139/cjm-2014-0668.
- Ullah S, M Asema and B Asghari (2014) Effect of PGPR on growth and performance of *Zea mays*. *Research Journal of Agriculture and Environmental Management* 2: 434-447.
- Verma JP, J Yadav, KN Tiwari, A Kumar (2013) Effect of indigenous Mesorhizobium spp. and plant growth promoting rhizobacteria on yields and nutrients uptake of chickpea (*Cicer arietinum* L.) under sustainable agriculture. *Ecol. Eng* 51: 282-286. 10.1016/j.ecoleng.2012.12.022.
- Wang J, R Li, H Zhang, G Wei and Z Li (2020) Beneficial bacteria activate nutrients and promote wheat growth under conditions of reduced fertilizer application. *BMC Microbiol* 20. <https://doi.org/10.1186/s12866-020-1708-z>.
- Wu SC, ZH Cao, ZG Li, KC Cheung and MH Wong (2005) Effects of Bio-fertilizer Containing N-Fixer, P and K Solubilizers and AM Fungi on Maize Growth: A Greenhouse Trial. *Geoderma* 125: 155-166. DOI: 10.1016/j.geoderma.2004.07.003.
- Yadav A, S Dhull, A Sehrawat and S Suneja (2017) Growth, survival and shelf life enhancement of phosphate solubilizing bacterial liquid inoculants formulations with polymeric additives. *Bioscan* 12:113-116.
- Yadav JJP Verma and KN Tiwari (2010) Effect of plant growth promoting rhizobacteria on seed germination and plant growth chickpea (*Cicer arietinum* L.) under in vitro conditions. *Biological Forum* 2: 15-18.
- Yazdani M, AM Bahmanyar, H Pirdashti, AM Esmaili (2009) Effect of phosphate solubilization microorganisms (PSM) and plant growth promoting rhizobacteria (PGPR) on yield and yield components of corn (*Zea mays* L.). *World Acad. Sci. Eng. Technol.* 49: 90-92. DOI: 10.5281/zenodo.1080014.
- Zaballa JI, R Golluscio and CM Ribaudo (2020) Effect of the Phosphorus-Solubilizing Bacterium *Enterobacter Ludwigii* on Barley Growth Promotion. *American Scientific Research Journal for Engineering, Technology, and Sciences (ASRJETS)* 63: 2313-4402.